


fertility 
associates



**Sperm microinjection
(ICSI) and surgical sperm
retrieval (SSR)**


Sperm microinjection (ICSI)

Sperm microinjection, commonly known as ICSI, means IVF can be used in almost any situation where there is male infertility, so long as there are some moving sperm. Even when sperm are not present in the ejaculate, they can often be obtained from a biopsy of the testis in a procedure known as surgical sperm retrieval (SSR).


Fertilisation of the egg by the sperm involves a complex sequence of events, initially controlled by the sperm. ICSI (intracytoplasmic sperm injection) bypasses the work the sperm usually has to do during fertilisation. During the ICSI procedure the embryologist captures sperm one at a time using a fine glass needle under a microscope and injects the sperm deep into the egg. **You can watch it here.**

ICSI is useful because in most men the genetic make-up of sperm is normal even though the sperm might be abnormally shaped, unable to move very well, or unable to undertake the steps of normal fertilisation.

Image of an egg during ICSI injection



The testis and epididymis, and ways of retrieving sperm



- A Site of PESA Epididymis (where sperm are stored and matured)
- B Vas deferens
- C Site of TESA, TESE or MESA Seminiferous tubules (where sperm are made)
- D Rete testis (which connects the seminiferous tubules)

Some men do not have any sperm in the ejaculate but still produce sperm in their testes. In this case, sperm can usually be obtained using Surgical Sperm Retrieval (SSR).

Sperm are produced in the seminiferous tubules in the testes, pass into the epididymis and then into the Vas deferens (see illustration).

It is the Vas deferens which is cut or blocked during a vasectomy – hence the name of the operation.

If a man has a blockage in the Vas deferens or in the epididymis outflow tract – from infection, vasectomy or even congenital absence of the vas – this is called 'obstructive azoospermia'. It is nearly always possible to retrieve sperm from the epididymis or testis from these men.

Other men sometimes have sperm production in the testis but the number

produced is too few for them to appear in the semen: this is called 'nonobstructive' azoospermia. In about 40% of men with nonobstructive azoospermia sperm our embryologists can sometimes find sperm in the ejaculate even when a pathology laboratory reports that there are no sperm present. We usually recommend at least two semen samples analysed by our by our embryologists before people think of moving on to SSR.

Usually the two types of azoospermia can be distinguished by physical examination and a blood test for the hormone FSH. Occasionally a testicular biopsy can help make a diagnosis. Because a testicular biopsy carries some risks, whether to have a biopsy or not needs careful discussion with your doctor.

Types of SSR

There are four types of SSR named after the techniques used:

- 1 TESA** (Testicular sperm aspiration) refers to passing a fine needle into the testes to withdraw seminiferous tubules from several areas within the testis.
- 2 TESE** (Testicular sperm extraction) refers to taking a larger amount of tissue from the testis through a cut in the skin.
- 3 PESA** (Percutaneous epididymal sperm aspiration) refers to passing a fine needle through the skin into the epididymis.
- 4 Microsurgical SSR** This is an operation done under general anaesthetic. The testis is removed from the scrotum and seminiferous tubules are examined under an operating microscope. This method reduces the risk of bleeding within the testis, and may increase the chance of finding sperm with nonobstructive azoospermia by about 10%. It usually only used for men with nonobstructive azoospermia, particularly if the testes are small. The procedure is more expensive than the other methods described below.

Local anaesthetic usually gives sufficient pain relief for PESA, TESA and TESE procedures.

PESA, TESA and TESE are usually done well before the start of an IVF cycle. If sperm are found, they can be frozen and then thawed later for use on the day of egg collection.

The number of straws needed to be thawed on the day of ICSI treatment will depend on the number and quality of the frozen sperm. Sometimes more straws have to be thawed than expected.

Risks and side effects of SSR

Nausea and loss of memory are common side effects of sedative and narcotic drugs used during SSR.

If analgesic drugs are used during biopsy they will affect your ability to drive safely, so you need to arrange transport home from the clinic. You cannot drive or use machinery during the next 24 hours, and we advise that someone remains with you for six hours after the procedure.

The drugs used for testicular biopsy can reduce the amount of air you breathe and thus the oxygen in your blood. If your oxygen saturation falls too low the doctor will stop the procedure and give oxygen. Very rarely you may require emergency drugs.

Bleeding and infection are possible complications of PESA, TESA and TESE, although they are rare. It is not uncommon to feel discomfort for several days, especially if TESE has been performed, and good scrotal support and Panadol is advised. If pain persists or recurs two or three days later, contact the clinic.

TESA and TESE are may cause inflammation in the testis that could reduce future sperm production. Up to 80% of men having TESE will have inflammation or collection of blood (called a haematoma) at the site of the biopsy. Complete loss of blood supply and atrophy of the testis has also been reported after a TESE procedure (rare).

A repeated TESE procedure is more likely to be successful if it is done at least six months after the last procedure, to allow the testis time to recover from temporary damage. Similar complications on a smaller scale may also occur after TESA.